



Respiratory Syncytial Virus


Respiratory syncytial (RES-pri-tor-ee sin-SISH-shul VYE-ris) *virus*, or RSV, is a common illness of the lungs and breathing passages affecting high-risk populations, especially babies born prematurely who are less than 2 years of age.


- By age 2, most children have had or been exposed to RSV.
- In healthy children, RSV is like a cold. It can be more serious for those in high-risk groups, such as
 - children more than 4 weeks premature at birth
 - twins or other multiple births
 - children younger than 2 years born with lung or heart disease
 - children who weighed less than 5.5 pounds at birth
 - school-aged brothers and sisters
 - children who attend day care

- children who live in a crowded home environment
- children with a family history of asthma
- children who are exposed to tobacco smoke and other air pollutants.

RSV can be dangerous for premature babies because premature babies, even those who did not require oxygen or positive pressure ventilation in the neonatal intensive care unit, do not have fully developed airways. In addition, they may not have received antibodies to help fight off RSV and other viruses because they were born early.

It is important that parents understand prior to discharge that it is essential during this first year of life to take specific precautions to decrease their baby's exposure to RSV. Precautions include continued proper hand washing and limiting their baby's exposure to crowds and school-aged





children. Healthcare professionals should discuss day care arrangements with parents and other intended caregivers. For infants with severe chronic lung disease or heart defects, traditional day care may not be the best alternative. The healthcare professional can help parents explore other options (e.g., private day care).

RSV infections usually begin in the fall and end late in the spring. Unfortunately, there also are several strains of RSV, so even if a baby has had it once during RSV season,

there is still a chance he or she can get it again. We each will have RSV several times in our lifetime.

RSV is very contagious. It can be spread through the air when a person coughs or sneezes. RSV also can spread by touching an object that has the virus on it. In fact, the virus can live on countertops, doorknobs, hands, and clothing for up to 7 hours. Hand washing and proper cleaning are the best ways to help prevent the spread of RSV. Synagis (palivizumab) is an immunoglobulin injection that helps prevent RSV. Synagis is not a vaccine, but it helps to prevent or lessen the symptoms should the infant contract RSV. The National Perinatal Association and the American Academy of Pediatrics each offer dosing guidelines (see links in bibliography below). Follow the dosing guidelines of your institution.

Bibliography

- American Academy of Pediatrics. (2009). Policy statement--modified recommendations for use of palivizumab for prevention of respiratory syncytial virus infections. *Pediatrics*, 124(6), 1694-1701. Retrieved from <http://pediatrics.aappublications.org/content/124/6/1694.full.pdf+html?sid=d220e4d6-0e9d-43a0-97b4-46604b1bfb46>
- Bracht, M., Heffer, M., & O'Brien, K. (2005). Preventing respiratory syncytial virus (RSV) infection. *Advances in Neonatal Care*, 5(1), 50-51.
- Centers for Disease Control and Prevention. (2013). Respiratory syncytial virus infection (RSV). Retrieved from www.cdc.gov/rsv
- National Perinatal Association. (2012). Respiratory syncytial virus (RSV) prevention 2012: Position statement. Retrieved from www.nationalperinatal.org/advocacy/pdf/Respiratory-Syncytial-Virus-Prevention-2012.pdf



Respiratory Syncytial Virus: Information for Parents

Respiratory syncytial (RES-pri-tor-ee sin-SISH-shul VYE-ris) *virus*, or RSV, is a virus that can make your baby sick. There are times of the year when your baby has more chance of being around people with the virus. This is usually from fall through spring (October through March). RSV can cause a mild cold, but it can also lead to a more serious sickness and a stay in the hospital. Babies born early and children younger than age 2 have the greatest risk for RSV. This is because babies born early, even those who did not need any help breathing while they were in the neonatal intensive care unit, do not have fully formed lungs. Also, their bodies may not be able to fight RSV and other viruses.

Symptoms of RSV

RSV usually causes a mild cold with a runny nose and fever. However, RSV also can make your baby very sick. Call your baby's provider right away if your baby has any of these symptoms:

- cough that does not go away, gets worse, or produces yellow, green, or grey mucus
- wheezing (a high-pitched whistling sound when breathing)

- trouble breathing or breathing faster than usual
- blue color on the lips or around the mouth
- high fever
- thick nasal discharge that is yellow, green, or grey.

There are ways to protect your baby. RSV is very easy to spread through coughing and sneezing. The virus can live on countertops, doorknobs, hands, and clothing for up to 7 hours. During RSV season, wash your hands well and avoid crowded places (like malls and churches) and school-age children. Speak with your baby's provider about your day care plans. You might have to tell other people what RSV is and how dangerous it can be for your baby. Your baby's provider can help with this by giving you some handouts to help you discuss RSV with friends and family.

Hand washing and keeping your baby away from sick people is a good way to protect your baby. There is also a medication that can help lower your baby's risk of getting an infection.